



How to Use AzaSite®



General patient information about AzaSite®

Do not allow the applicator tip to touch the eye, fingers, or other sources.

Stop use and contact a physician if any signs of an allergic reaction occur.

Although it is common to feel better soon after starting therapy, AzaSite® should be taken exactly as directed. Skipping doses or not completing the full course of therapy may

- Decrease the effectiveness of the current treatment
- Increase the likelihood that bacteria will develop resistance and will not be treatable by AzaSite® or other antibacterial drugs in the future

Do not wear contact lenses if you have signs or symptoms of bacterial conjunctivitis.


Wash hands thoroughly before using AzaSite®.

It is important to follow the steps below for proper administration.

Please follow steps 1 and 2 the *first* time you use a new bottle of AzaSite®:




1. Remove the white tamper-evident cap by turning it in a clockwise direction. Discard the white cap.




2. Holding the bottle upright, remove the tan cap by turning it in a counterclockwise direction. This will allow any excess pressure in the bottle to be released. Replace the tan cap on the bottle and close securely.


Please follow steps 3 through 6 *each* time you use AzaSite®:



3. Turn the closed bottle upside down.



4. Shake hand firmly before each use to move the contents into the tip of the bottle.



5. Keeping the bottle upside down, remove the tan cap.



6. Tilt head back. With the bottle inverted, gently squeeze the bottle to instill 1 drop into each affected eye. Replace the tan cap back on the bottle and close securely after each administration for proper storage.

AzaSite® (azithromycin ophthalmic solution) 1% is indicated for the treatment of bacterial conjunctivitis in adults and children 1 year of age and older. Recommended dosing for AzaSite® is 1 drop twice a day for 2 days, followed by 1 drop once a day for the next 5 days, for a total of 9 drops per affected eye.

Important Safety Information: For use as an eyedrop only. In clinical trials, the most common ocular adverse event was eye irritation, which occurred in 1% to 2% of patients.

Please see accompanying full Prescribing Information.

Manufactured for: Inspire Pharmaceuticals, Inc., a subsidiary of
MERCK & CO., INC., Whitehouse Station, NJ 08889, USA

AzaSite is a registered trademark of InSite Vision Inc.

Copyright © 2011 Inspire Pharmaceuticals, Inc., a subsidiary of Merck & Co., Inc.

All rights reserved.

AzaSITE
(azithromycin ophthalmic solution) 1%

