

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use DELSTRIGO safely and effectively. See full prescribing information for DELSTRIGO.

DELSTRIGO® (doravirine, lamivudine, and tenofovir disoproxil fumarate) tablets, for oral use
Initial U.S. Approval: 2018

WARNING: POSTTREATMENT ACUTE EXACERBATION OF HEPATITIS B

See full prescribing information for complete boxed warning.

Severe acute exacerbations of hepatitis B (HBV) have been reported in people with concomitant HIV-1 and HBV who have discontinued lamivudine or tenofovir disoproxil fumarate (TDF), two of the components of DELSTRIGO. Closely monitor hepatic function in these patients. If appropriate, initiation of anti-hepatitis B therapy may be warranted. (5.2)

RECENT MAJOR CHANGES

Warnings and Precautions (5.1)	11/2024
Warnings and Precautions (5.5)	06/2025

INDICATIONS AND USAGE

DELSTRIGO is a three-drug combination of doravirine (a non-nucleoside reverse transcriptase inhibitor [NNRTI]), lamivudine, and tenofovir disoproxil fumarate (both nucleoside analogue reverse transcriptase inhibitors) and is indicated as a complete regimen for the treatment of HIV-1 infection in adults and pediatric patients weighing at least 35 kg:

- with no antiretroviral treatment history, **OR**
- to replace the current antiretroviral regimen in those who are virologically-suppressed (HIV-1 RNA less than 50 copies per mL) on a stable antiretroviral regimen with no history of treatment failure and no known substitutions associated with resistance to the individual components of DELSTRIGO. (1)

DOSAGE AND ADMINISTRATION

- Testing: Prior to or when initiating DELSTRIGO, test for HBV infection. Prior to or when initiating DELSTRIGO, and during treatment with DELSTRIGO, on a clinically appropriate schedule, assess serum creatinine, estimated creatinine clearance, urine glucose and urine protein in all patients. In patients with chronic kidney disease, also assess serum phosphorus. (2.1)
- Recommended dosage: One tablet taken orally once daily with or without food in adults and pediatric patients weighing at least 35 kg. (2.2)
- Renal impairment: Not recommended in patients with estimated creatinine clearance below 50 mL per minute. (2.3)
- Dosage adjustment with rifabutin: Take one tablet of DELSTRIGO once daily, followed by one tablet of doravirine 100 mg (PIFELTRO) approximately 12 hours after the dose of DELSTRIGO. (2.4)

FULL PRESCRIBING INFORMATION: CONTENTS*

WARNING: POSTTREATMENT ACUTE EXACERBATION OF HEPATITIS B

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Testing When Initiating and During Treatment with DELSTRIGO
- 2.2 Recommended Dosage
- 2.3 Renal Impairment
- 2.4 Dosage Adjustment with Rifabutin

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Severe Skin Reactions
- 5.2 Severe Acute Exacerbation of Hepatitis B in people with concomitant HIV-1 and HBV
- 5.3 New Onset or Worsening Renal Impairment
- 5.4 Risk of Adverse Reactions or Loss of Virologic Response Due to Drug Interactions
- 5.5 Bone Loss and Mineralization Defects

DOSAGE FORMS AND STRENGTHS

- Tablets: 100 mg of doravirine, 300 mg of lamivudine, and 300 mg of tenofovir disoproxil fumarate. (3)

CONTRAINDICATIONS

- DELSTRIGO is contraindicated when co-administered with drugs that are strong cytochrome P450 (CYP)3A enzyme inducers as significant decreases in doravirine plasma concentrations may occur, which may decrease the effectiveness of DELSTRIGO. (4)
- DELSTRIGO is contraindicated in patients with a previous hypersensitivity reaction to lamivudine.

WARNINGS AND PRECAUTIONS

- Severe skin reactions, including Stevens-Johnson syndrome (SJS)/toxic epidermal necrolysis (TEN), have been reported during the postmarketing experience with doravirine-containing regimens. Discontinue DELSTRIGO, and other medications known to be associated with severe skin reactions, immediately if a painful rash with mucosal involvement or a progressive severe rash develops, and closely monitor clinical status. (5.1)
- New onset or worsening renal impairment: Prior to or when initiating DELSTRIGO, and during treatment with DELSTRIGO, on a clinically appropriate schedule, assess serum creatinine, estimated creatinine clearance, urine glucose, and urine protein in all patients. Avoid administering DELSTRIGO with concurrent or recent use of nephrotoxic drugs. (5.3)
- Bone loss and mineralization defects: Consider monitoring BMD in patients with a history of pathologic fracture or other risk factors of osteoporosis or bone loss. (5.5)
- Monitor for Immune Reconstitution Syndrome. (5.6)

ADVERSE REACTIONS

Most common adverse reactions (incidence greater than or equal to 5%, all grades) are dizziness, nausea, and abnormal dreams. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Merck Sharp & Dohme LLC at 1-877-888-4231 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- Because DELSTRIGO is a complete regimen, co-administration with other antiretroviral medications for treatment of HIV-1 infection is not recommended. (7.1)
- Consult the full prescribing information prior to and during treatment for important potential drug-drug interactions. (4, 5.4, 7)

USE IN SPECIFIC POPULATIONS

- Pediatrics: Not recommended for patients weighing less than 35 kg. (8.4)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 06/2025

- 5.6 Immune Reconstitution Syndrome
- 6 ADVERSE REACTIONS
 - 6.1 Clinical Trials Experience
 - 6.2 Postmarketing Experience
- 7 DRUG INTERACTIONS
 - 7.1 Concomitant Use with Other Antiretroviral Medications
 - 7.2 Effect of Other Drugs on DELSTRIGO
 - 7.3 Effect of DELSTRIGO on Other Drugs
- 8 USE IN SPECIFIC POPULATIONS
 - 8.1 Pregnancy
 - 8.2 Lactation
 - 8.4 Pediatric Use
 - 8.5 Geriatric Use
 - 8.6 Renal Impairment
 - 8.7 Hepatic Impairment
- 10 OVERDOSAGE
- 11 DESCRIPTION
- 12 CLINICAL PHARMACOLOGY
 - 12.1 Mechanism of Action
 - 12.2 Pharmacodynamics

12.3 Pharmacokinetics

12.4 Microbiology

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

14.1 Clinical Trial Results in Adults with No Antiretroviral Treatment History

14.2 Clinical Trial Results in Virologically-Suppressed Adults

14.3 Clinical Trial Results in Pediatric Participants

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

*Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

WARNING: POSTTREATMENT ACUTE EXACERBATION OF HEPATITIS B

Severe acute exacerbations of hepatitis B (HBV) have been reported in people with concomitant HIV-1 and HBV who have discontinued lamivudine or tenofovir disoproxil fumarate (TDF), which are components of DELSTRIGO. Hepatic function should be monitored closely with both clinical and laboratory follow-up for at least several months in patients who are coinfecting with HIV-1 and HBV and discontinue DELSTRIGO. If appropriate, initiation of anti-hepatitis B therapy may be warranted [see *Warnings and Precautions* (5.2)].

1 INDICATIONS AND USAGE

DELSTRIGO® is indicated as a complete regimen for the treatment of HIV-1 infection in adults and pediatric patients weighing at least 35 kg:

- with no prior antiretroviral treatment history, **OR**
- to replace the current antiretroviral regimen in those who are virologically-suppressed (HIV-1 RNA less than 50 copies per mL) on a stable antiretroviral regimen with no history of treatment failure and no known substitutions associated with resistance to the individual components of DELSTRIGO [see *Clinical Studies* (14)].

2 DOSAGE AND ADMINISTRATION

2.1 Testing When Initiating and During Treatment with DELSTRIGO

Prior to or when initiating DELSTRIGO, test patients for HBV infection [see *Warnings and Precautions* (5.2)].

Prior to or when initiating DELSTRIGO, and during treatment with DELSTRIGO, on a clinically appropriate schedule, assess serum creatinine, estimated creatinine clearance, urine glucose, and urine protein in all patients. In patients with chronic kidney disease, also assess serum phosphorus [see *Warnings and Precautions* (5.3)].

2.2 Recommended Dosage

DELSTRIGO is a fixed-dose combination product containing 100 mg of doravirine (DOR), 300 mg of lamivudine (3TC), and 300 mg of TDF. The recommended dosage of DELSTRIGO in adults and pediatric patients weighing at least 35 kg is one tablet taken orally once daily with or without food [see *Clinical Pharmacology* (12.3)].

2.3 Renal Impairment

Because DELSTRIGO is a fixed-dose combination tablet and the dosage of lamivudine and TDF cannot be adjusted, DELSTRIGO is not recommended in patients with estimated creatinine clearance less than 50 mL/min [see *Warnings and Precautions* (5.3) and *Use in Specific Populations* (8.6)].

2.4 Dosage Adjustment with Rifabutin

If DELSTRIGO is co-administered with rifabutin, take one tablet of DELSTRIGO once daily, followed by one tablet of doravirine 100 mg (PIFELTRO) approximately 12 hours after the dose of DELSTRIGO for the duration of rifabutin co-administration [see *Drug Interactions* (7.2) and *Clinical Pharmacology* (12.3)].

3 DOSAGE FORMS AND STRENGTHS

DELSTRIGO film-coated tablets are yellow, oval-shaped tablets, debossed with the corporate logo and 776 on one side and plain on the other side. Each tablet contains 100 mg doravirine, 300 mg lamivudine, and 300 mg tenofovir disoproxil fumarate (equivalent to 245 mg of tenofovir disoproxil).

4 CONTRAINDICATIONS

- DELSTRIGO is contraindicated when co-administered with drugs that are strong cytochrome P450 (CYP)3A enzyme inducers as significant decreases in doravirine plasma concentrations may occur, which may decrease the effectiveness of DELSTRIGO [see *Warnings and Precautions* (5.4), *Drug Interactions* (7.2), and *Clinical Pharmacology* (12.3)]. These drugs include, but are not limited to, the following:
 - the anticonvulsants carbamazepine, oxcarbazepine, phenobarbital, phenytoin
 - the androgen receptor inhibitor enzalutamide
 - the antimycobacterials rifampin, rifapentine
 - the cytotoxic agent mitotane
 - St. John's wort (*Hypericum perforatum*)
- DELSTRIGO is contraindicated in patients with a previous hypersensitivity reaction to lamivudine.

5 WARNINGS AND PRECAUTIONS

5.1 Severe Skin Reactions

Severe skin reactions, including Stevens-Johnson syndrome (SJS)/toxic epidermal necrolysis (TEN), have been reported during the postmarketing experience with doravirine-containing regimens [see *Adverse Reactions* (6.2)]. Discontinue DELSTRIGO, and other medications known to be associated with severe skin reactions, immediately if a painful rash with mucosal involvement or a progressive severe rash develops. Clinical status should be closely monitored, and appropriate therapy should be initiated.

5.2 Severe Acute Exacerbation of Hepatitis B in People with Concomitant HIV-1 and HBV

All patients with HIV-1 should be tested for the presence of HBV before initiating antiretroviral therapy.

Severe acute exacerbations of hepatitis B (e.g., liver decompensated and liver failure) have been reported in people with concomitant HIV-1 and HBV who have discontinued products containing lamivudine and/or TDF, and may occur with discontinuation of DELSTRIGO. Patients who are coinfectd with HIV-1 and HBV who discontinue DELSTRIGO should be closely monitored with both clinical and laboratory follow-up for at least several months after stopping treatment with DELSTRIGO. If appropriate, initiation of anti-hepatitis B therapy may be warranted, especially in patients with advanced liver disease or cirrhosis, since post-treatment exacerbation of hepatitis may lead to hepatic decompensation and liver failure.

5.3 New Onset or Worsening Renal Impairment

Renal impairment, including cases of acute renal failure and Fanconi syndrome (renal tubular injury with severe hypophosphatemia), has been reported with the use of TDF, a component of DELSTRIGO.

DELSTRIGO should be avoided with concurrent or recent use of a nephrotoxic agent (e.g., high-dose or multiple nonsteroidal anti-inflammatory drugs [NSAIDs]) [see *Drug Interactions* (7.1)]. Cases of acute renal failure after initiation of high-dose or multiple NSAIDs have been reported in patients living with HIV with risk factors for renal dysfunction who appeared stable on TDF. Some patients required hospitalization and renal replacement therapy. Alternatives to NSAIDs should be considered, if needed, in patients at risk for renal dysfunction.

Persistent or worsening bone pain, pain in extremities, fractures, and/or muscular pain or weakness may be manifestations of proximal renal tubulopathy and should prompt an evaluation of renal function in at-risk patients.

Prior to or when initiating DELSTRIGO, and during treatment with DELSTRIGO, on a clinically appropriate schedule, assess serum creatinine, estimated creatinine clearance, urine glucose and urine protein in all patients. In patients with chronic kidney disease, also assess serum phosphorus. Discontinue DELSTRIGO in patients who develop clinically significant decreases in renal function or evidence of Fanconi syndrome.

The lamivudine and TDF components of DELSTRIGO are primarily excreted by the kidney. Discontinue DELSTRIGO if estimated creatinine clearance declines below 50 mL/min as dose interval adjustment required for lamivudine and TDF cannot be achieved with the fixed-dose combination tablet [see *Use in Specific Populations* (8.6)].

5.4 Risk of Adverse Reactions or Loss of Virologic Response Due to Drug Interactions

The concomitant use of DELSTRIGO and certain other drugs may result in known or potentially significant drug interactions, some of which may lead to [see *Dosage and Administration* (2.4), *Contraindications* (4), and *Drug Interactions* (7.2)]:

- Loss of therapeutic effect of DELSTRIGO and possible development of resistance.
- Possible clinically significant adverse reactions from greater exposures of a component of DELSTRIGO.

See Table 6 for steps to prevent or manage these possible and known significant drug interactions, including dosing recommendations. Consider the potential for drug interactions prior to and during DELSTRIGO therapy, review concomitant medications during DELSTRIGO therapy, and monitor for adverse reactions.

5.5 Bone Loss and Mineralization Defects

Bone Mineral Density

In clinical trials in adults living with HIV, TDF (a component of DELSTRIGO) was associated with slightly greater decreases in bone mineral density (BMD) and increases in biochemical markers of bone metabolism, suggesting increased bone turnover relative to comparators. Serum parathyroid hormone levels and 1,25 Vitamin D levels were also higher in participants receiving TDF.

Clinical trials evaluating TDF in pediatric participants were conducted. Under normal circumstances, BMD increases rapidly in pediatric patients. In participants 2 years to less than 18 years of age living with HIV, bone effects were similar to those observed in adult participants and suggest increased bone turnover. Total body BMD gain was less in the TDF-treated pediatric participants living with HIV as compared to the control groups. Similar trends were observed in chronic HBV-infected pediatric participants 2 years to less than 18 years of age. In all pediatric trials, normal skeletal growth (height) was not affected for the duration of the clinical trials.

The effects of TDF-associated changes in BMD and biochemical markers on long-term bone health and future fracture risk in adults and pediatric participants 2 years and older are unknown. The long-term effect of lower spine and total body BMD on skeletal growth in pediatric patients, and in particular, the effects of long-duration exposure in younger children are unknown.

Assessment of BMD should be considered for adult and pediatric patients living with HIV who have a history of pathologic bone fracture or other risk factors for osteoporosis or bone loss. Although the effect of supplementation with calcium and vitamin D was not studied, such supplementation may be beneficial in all patients. If bone abnormalities are suspected, then appropriate consultation should be obtained.

Mineralization Defects

Cases of osteomalacia associated with proximal renal tubulopathy, manifested as bone pain or pain in extremities and which may contribute to fractures, have been reported in association with the use of TDF [see *Adverse Reactions* (6.2)]. Arthralgias and muscle pain or weakness have also been reported in cases of proximal renal tubulopathy. Hypophosphatemia and osteomalacia secondary to proximal renal tubulopathy should be considered in patients at risk of renal dysfunction who present with persistent or worsening bone or muscle symptoms while receiving products containing TDF [see *Warnings and Precautions* (5.3)].

5.6 Immune Reconstitution Syndrome

Immune reconstitution syndrome has been reported in patients treated with combination antiretroviral therapy. During the initial phase of combination antiretroviral treatment, patients whose immune system responds may develop an inflammatory response to indolent or residual opportunistic infections (such as *Mycobacterium avium* infection, cytomegalovirus, *Pneumocystis jirovecii* pneumonia (PCP), or tuberculosis), which may necessitate further evaluation and treatment.

Autoimmune disorders (such as Graves' disease, polymyositis, Guillain-Barré syndrome, and autoimmune hepatitis) have also been reported to occur in the setting of immune reconstitution; however, the time to onset is more variable and can occur many months after initiation of treatment.

6 ADVERSE REACTIONS

The following adverse reactions are discussed in other sections of the labeling:

- Severe Acute Exacerbation of Hepatitis B in people with concomitant HIV-1 and HBV [see *Warnings and Precautions* (5.2)]
- New Onset or Worsening Renal Impairment [see *Warnings and Precautions* (5.3)]
- Bone Loss and Mineralization Defects [see *Warnings and Precautions* (5.5)]
- Immune Reconstitution Syndrome [see *Warnings and Precautions* (5.6)]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Adverse Reactions in Adults with No Antiretroviral Treatment History

The safety assessment of DELSTRIGO is based on Week 96 data from two Phase 3, randomized, international, multicenter, double-blind, active-controlled trials. A total of 747 participants received doravirine either as the single entity in combination with other antiretroviral drugs as background regimens (n=383) or as the fixed-dose DELSTRIGO (n=364), and a total of 747 participants were randomized to control arms.

In DRIVE-AHEAD (Protocol 021), 728 adult participants received either DELSTRIGO (n=364) or EFV/FTC/TDF once daily (n=364). By Week 96, 3% in the DELSTRIGO group and 7% in the EFV/FTC/TDF group had adverse events leading to discontinuation of study medication.

Adverse reactions reported in greater than or equal to 5% of participants in any treatment group in DRIVE-AHEAD are presented in Table 1.

Table 1: Adverse Reactions* (All Grades) Reported in ≥5%[†] of Participants in Any Treatment Group in Adults with No Antiretroviral Treatment History in DRIVE-AHEAD (Week 96)

	DELSTRIGO Once Daily N=364	EFV/FTC/TDF Once Daily N=364
Dizziness	7%	32%
Nausea	5%	7%
Abnormal Dreams	5%	10%
Headache	4%	5%

Insomnia	4%	5%
Diarrhea	4%	6%
Somnolence	3%	7%
Rash [‡]	2%	12%
<p>*Frequencies of adverse reactions are based on all adverse events attributed to trial drugs by the investigator.</p> <p>[†]No adverse reactions of Grade 2 or higher (moderate or severe) occurred in ≥2% of participants treated with DELSTRIGO.</p> <p>[‡]Rash: includes rash, rash erythematous, rash generalized, rash macular, rash maculopapular, rash papular, rash pruritic.</p>		

The majority (66%) of adverse reactions associated with DELSTRIGO occurred at severity Grade 1 (mild).

Neuropsychiatric Adverse Events

For DRIVE-AHEAD, the analysis of participants with neuropsychiatric adverse events by Week 48 is presented in Table 2. The proportion of participants who reported one or more neuropsychiatric adverse events was 24% and 57% in the DELSTRIGO and EFV/FTC/TDF groups, respectively.

A statistically significantly lower proportion of DELSTRIGO-treated participants compared to EFV/FTC/TDF-treated participants reported neuropsychiatric adverse events by Week 48 in the three pre-specified categories of dizziness, sleep disorders and disturbances, and altered sensorium.

Table 2: DRIVE-AHEAD - Analysis of Participants with Neuropsychiatric Adverse Events* (Week 48)

	DELSTRIGO Once Daily N=364	EFV/FTC/TDF Once Daily N=364	Treatment Difference (DELSTRIGO - EFV/FTC/TDF) Estimate (95% CI)[†]
Sleep disorders and disturbances [‡]	12%	26%	-13.5 (-19.1, -7.9)
Dizziness	9%	37%	-28.3 (-34.0, -22.5)
Altered sensorium [§]	4%	8%	-3.8 (-7.6, -0.3)
<p>*All causality and all grade events were included in the analysis.</p> <p>[†]The 95% CIs were calculated using Miettinen and Nurminen's method. Categories pre-specified for statistical testing were dizziness (p <0.001), sleep disorders and disturbances (p <0.001), and altered sensorium (p=0.033).</p> <p>[‡]Predefined using MedDRA preferred terms including: abnormal dreams, hypsomnias, initial insomnia, insomnia, nightmare, sleep disorder, somnambulism.</p> <p>[§]Predefined using MedDRA preferred terms including: altered state of consciousness, lethargy, somnolence, syncope.</p>			

Neuropsychiatric adverse events in the pre-defined category of depression and suicide/self-injury were reported in 4% and 7% of participants, in the DELSTRIGO and EFV/FTC/TDF groups, respectively.

In DRIVE-AHEAD through 48 weeks of treatment, the majority of participants who reported neuropsychiatric adverse events reported events that were mild to moderate in severity (97% [83/86] and 96% [198/207], in the DELSTRIGO and EFV/FTC/TDF groups, respectively) and the majority of participants reported these events in the first 4 weeks of treatment (72% [62/86] in the DELSTRIGO group and 86% [177/207] in the EFV/FTC/TDF group).

Neuropsychiatric adverse events led to treatment discontinuation in 1% (2/364) and 1% (5/364) of participants in the DELSTRIGO and EFV/FTC/TDF groups, respectively. The proportion of participants who

reported neuropsychiatric adverse events through Week 4 was 17% (62/364) in the DELSTRIGO group and 49% (177/364) in the EFV/FTC/TDF group. At Week 48, the prevalence of neuropsychiatric adverse events was 12% (44/364) in the DELSTRIGO group and 22% (81/364) in the EFV/FTC/TDF group. At Week 96, the prevalence of neuropsychiatric adverse events was 13% (47/364) in the DELSTRIGO group and 23% (82/364) in the EFV/FTC/TDF group.

Laboratory Abnormalities

The percentages of participants with selected laboratory abnormalities (that represent a worsening from baseline) who were treated with DELSTRIGO or EFV/FTC/TDF in DRIVE-AHEAD are presented in Table 3.

Table 3: Selected Laboratory Abnormalities Reported in Adult Participants with No Antiretroviral Treatment History in DRIVE-AHEAD (Week 96)

Laboratory Parameter Preferred Term (Unit)/Limit	DELSTRIGO Once Daily N=364	EFV/FTC/TDF Once Daily N=364
Blood Chemistry		
Total bilirubin (mg/dL)		
1.1 - <1.6 x ULN	5%	0%
1.6 - <2.6 x ULN	2%	0%
≥2.6 x ULN	1%	<1%
Creatinine (mg/dL)		
>1.3 - 1.8 x ULN or Increase of >0.3 mg/dL above baseline	3%	2%
>1.8 x ULN or Increase of ≥1.5 x above baseline	3%	2%
Aspartate aminotransferase (IU/L)		
2.5 - <5.0 x ULN	3%	3%
≥5.0 x ULN	1%	4%
Alanine aminotransferase (IU/L)		
2.5 - <5.0 x ULN	4%	4%
≥5.0 x ULN	1%	3%
Alkaline phosphatase (IU/L)		
2.5 - <5.0 x ULN	<1%	1%
≥5.0 x ULN	0%	<1%
Lipase		
1.5 - <3.0 x ULN	6%	4%
≥3.0 x ULN	2%	3%
Creatine kinase (IU/L)		
6.0 - <10.0 x ULN	3%	3%
≥10.0 x ULN	4%	6%
Cholesterol, fasted (mg/dL)		
≥300 mg/dL	1%	<1%
LDL cholesterol, fasted (mg/dL)		
≥190 mg/dL	<1%	2%
Triglycerides, fasted (mg/dL)		
>500 mg/dL	1%	3%
Each participant is only counted once per parameter at the highest toxicity grade. Only participants with a baseline value and at least one on-treatment value for a given laboratory parameter are included. ULN = Upper limit of normal range.		

Change in Lipids from Baseline

For DRIVE-AHEAD, changes from baseline at Week 48 in LDL-cholesterol, non-HDL-cholesterol, total cholesterol, triglycerides, and HDL-cholesterol are shown in Table 4. Changes from baseline at Week 96 were similar to findings at Week 48.

The LDL and non-HDL comparisons were pre-specified and are summarized in Table 4. The differences were statistically significant, showing superiority of DELSTRIGO for both parameters. The clinical benefit of these findings has not been demonstrated.

Table 4: Mean Change from Baseline in Fasting Lipids in Adult Participants with No Antiretroviral Treatment History in DRIVE-AHEAD (Week 48)

Laboratory Parameter Preferred Term	DELSTRIGO Once Daily N=320		EFV/FTC/TDF Once Daily N=307		Difference Estimates (DELSTRIGO - EFV/FTC/TDF)
	Baseline	Change	Baseline	Change	Difference (95% CI)
LDL-Cholesterol (mg/dL)*	91.7	-2.1	91.3	8.3	-10.2 (-13.8, -6.7)
Non-HDL Cholesterol (mg/dL)*	114.7	-4.1	115.3	12.7	-16.9 (-20.8, -13.0)
Total Cholesterol (mg/dL)†	156.8	-2.2	156.8	21.1	-
Triglycerides (mg/dL)†	118.7	-12.0	122.6	21.6	-
HDL-Cholesterol (mg/dL)†	42.1	1.8	41.6	8.4	-
Participants on lipid-lowering agents at baseline were excluded from these analyses (DELSTRIGO n=15 and EFV/FTC/TDF n=10). Participants initiating a lipid-lowering agent post-baseline had their last fasted on-treatment value (prior to starting the agent) carried forward (DELSTRIGO n=3 and EFV/FTC/TDF n=8).					
*p-value for the pre-specified hypothesis testing for treatment difference was <0.0001.					
†Not pre-specified for hypothesis testing.					

Adverse Reactions in Virologically-Suppressed Adults

The safety of DELSTRIGO in virologically-suppressed adults was based on Week 48 data from 670 participants in the DRIVE-SHIFT trial (Protocol 024), a randomized, international, multicenter, open-label trial in which virologically-suppressed participants were switched from a baseline regimen consisting of two nucleoside reverse transcriptase inhibitors (NRTIs) in combination with a protease inhibitor (PI) plus either ritonavir or cobicistat, or elvitegravir plus cobicistat, or an NNRTI to DELSTRIGO. Overall, the safety profile in virologically-suppressed adult participants was similar to that in participants with no antiretroviral treatment history.

Laboratory Abnormalities

Serum ALT and AST Elevations: In the DRIVE-SHIFT trial, 22% and 16% of participants in the immediate switch group experienced ALT and AST elevations of greater than 1.25 X ULN, respectively, through 48 weeks on DELSTRIGO. For these ALT and AST elevations, no apparent time patterns with regard to time to onset relative to switch were observed. One percent of participants had ALT or AST elevations greater than 5 X ULN through 48 weeks on DELSTRIGO. The ALT and AST elevations were generally asymptomatic, and not associated with bilirubin elevations. In comparison, 4% and 4% of participants in the delayed switch group experienced ALT and AST elevations of greater than 1.25 X ULN through 24 weeks on their baseline regimen.

Change in Lipids from Baseline

Changes from baseline at Week 24 in LDL-cholesterol, non-HDL-cholesterol, total cholesterol, triglycerides, and HDL-cholesterol in participants on a PI plus ritonavir-based regimen at baseline are shown in Table 5. The LDL and non-HDL comparisons were pre-specified, and the differences were statistically significant, showing superiority for an immediate switch to DELSTRIGO for both parameters. The clinical benefit of these findings has not been demonstrated.

Table 5: Mean Change from Baseline in Fasting Lipids in Adult Virologically-Suppressed Participants on a PI plus Ritonavir-based Regimen at Baseline in DRIVE-SHIFT (Week 24)

Laboratory Parameter Preferred Term	DELSTRIGO (Week 0-24) Once Daily N=244		PI+ritonavir (Week 0-24) Once Daily N=124		Difference Estimates
	Baseline	Change	Baseline	Change	Difference (95% CI)
LDL-Cholesterol (mg/dL)*	108.7	-16.3	110.5	-2.6	-14.5 (-18.9, -10.1)
Non-HDL Cholesterol (mg/dL)*	138.6	-24.8	138.8	-2.1	-22.8 (-27.9, -17.7)
Total Cholesterol (mg/dL)†	188.5	-26.1	187.4	-0.2	-
Triglycerides (mg/dL)†	153.1	-44.4	151.4	-0.4	-
HDL-Cholesterol (mg/dL)†	50.0	-1.3	48.5	1.9	-
Participants on lipid-lowering agents at baseline were excluded from these analyses (DELSTRIGO n=26 and PI+ritonavir n=13). Participants initiating a lipid-lowering agent post-baseline had their last fasted on-treatment value (prior to starting the agent) carried forward (DELSTRIGO n=4 and PI+ritonavir n=2). *p-value for the pre-specified hypothesis testing for treatment difference was <0.0001. †Not pre-specified for hypothesis testing.					

Adverse Reactions in Pediatric Participants

The safety of DELSTRIGO was evaluated in 45 virologically-suppressed or treatment-naïve pediatric participants 12 to less than 18 years of age living with HIV, through Week 24 in an open-label trial (IMPAACT 2014 (Protocol 027)) [see *Clinical Studies* (14.3)]. The safety profile in pediatric participants was similar to that in adults. There were no serious or Grade 3 or 4 adverse reactions. No participants discontinued due to an adverse event.

6.2 Postmarketing Experience

The following adverse reactions have been identified during postmarketing experience in patients receiving doravirine-, lamivudine- or TDF-containing regimens. Because postmarketing reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Doravirine:

Skin and Subcutaneous Tissue Disorders: Stevens-Johnson syndrome (SJS)/toxic epidermal necrolysis (TEN)

Lamivudine:

Body as a Whole: redistribution/accumulation of body fat

Endocrine and Metabolic: hyperglycemia

General: Weakness

Hemic and Lymphatic: anemia (including pure red cell aplasia and severe anemias progressing on therapy)

Hepatic and Pancreatic: lactic acidosis and hepatic steatosis, posttreatment exacerbations of hepatitis B

Hypersensitivity: anaphylaxis, urticaria

Musculoskeletal: muscle weakness, CPK elevation, rhabdomyolysis

Skin: alopecia, pruritus

TDF:

Immune System Disorders: allergic reaction, including angioedema

Metabolism and Nutrition Disorders: lactic acidosis, hypokalemia, hypophosphatemia

Respiratory, Thoracic, and Mediastinal Disorders: dyspnea

Gastrointestinal Disorders: pancreatitis, increased amylase, abdominal pain

Hepatobiliary Disorders: hepatic steatosis, hepatitis, increased liver enzymes (most commonly AST, ALT gamma GT)

Skin and Subcutaneous Tissue Disorders: rash

Musculoskeletal and Connective Tissue Disorders: rhabdomyolysis, osteomalacia (manifested as bone pain and which may contribute to fractures), muscular weakness, myopathy

Renal and Urinary Disorders: acute renal failure, renal failure, acute tubular necrosis, Fanconi syndrome, proximal renal tubulopathy, interstitial nephritis (including acute cases), nephrogenic diabetes insipidus, renal insufficiency, increased creatinine, proteinuria, polyuria

General Disorders and Administration Site Conditions: asthenia

The following adverse reactions, listed under the body system headings above, may occur as a consequence of proximal renal tubulopathy: rhabdomyolysis, osteomalacia, hypokalemia, muscular weakness, myopathy, hypophosphatemia.

7 DRUG INTERACTIONS

7.1 Concomitant Use with Other Antiretroviral Medications

Because DELSTRIGO is a complete regimen for the treatment of HIV-1 infection, co-administration with other antiretroviral medications for treatment of HIV-1 infection is not recommended. Information regarding potential drug-drug interactions with other antiretroviral medications is not provided.

7.2 Effect of Other Drugs on DELSTRIGO

Co-administration of DELSTRIGO with a CYP3A inducer decreases doravirine plasma concentrations, which may reduce DELSTRIGO efficacy [see *Contraindications (4)*, *Warnings and Precautions (5.4)*, and *Clinical Pharmacology (12.3)*].

Co-administration of DELSTRIGO and drugs that are inhibitors of CYP3A may result in increased plasma concentrations of doravirine.

Table 6 shows the significant drug interactions with the components of DELSTRIGO. The drug interactions described are based on studies conducted with either DELSTRIGO or the components of DELSTRIGO as individual agents.

Table 6: Drug Interactions with DELSTRIGO*

Concomitant Drug Class: Drug Name	Effect on Concentration	Clinical Comment
Androgen Receptors		
enzalutamide	↓ doravirine	Co-administration is contraindicated with enzalutamide. At least a 4-week cessation period is recommended prior to initiation of DELSTRIGO.
Anticonvulsants		
carbamazepine oxcarbazepine phenobarbital phenytoin	↓ doravirine	Co-administration is contraindicated with these anticonvulsants. At least a 4-week cessation period is recommended prior to initiation of DELSTRIGO.
Antimycobacterials		
rifampin† rifapentine	↓ doravirine	Co-administration is contraindicated with rifampin or rifapentine. At least a 4-week cessation period is recommended prior to initiation of DELSTRIGO.
rifabutin†	↓ doravirine	If DELSTRIGO is co-administered with rifabutin, one tablet of doravirine (PIFELTRO) should be taken approximately 12 hours after the dose of DELSTRIGO [see <i>Dosage and Administration</i> (2.4)].
Cytotoxic Agents		
mitotane	↓ doravirine	Co-administration is contraindicated with mitotane. At least a 4-week cessation period is recommended prior to initiation of DELSTRIGO.
Hepatitis C Antiviral Agents		
ledipasvir/sofosbuvir sofosbuvir/velpatasvir	↑ tenofovir	Monitor for adverse reactions associated with TDF.
Herbal Products		
St. John's wort	↓ doravirine	Co-administration is contraindicated with St. John's wort. At least a 4-week cessation period is recommended prior to initiation of DELSTRIGO.
Other Agents		
sorbitol	↓ lamivudine	Co-administration of single doses of lamivudine and sorbitol resulted in a sorbitol dose-dependent reduction in lamivudine exposures. When possible, avoid use of sorbitol-containing medicines with lamivudine-containing medicines.
↑ = increase, ↓ = decrease *This table is not all-inclusive †The interaction between doravirine and the concomitant drug was evaluated in a clinical study. All other drug-drug interactions shown are anticipated based on the known metabolic and elimination pathways.		

Co-administration of DELSTRIGO with drugs that reduce renal function or compete for active tubular secretion may increase serum concentrations of lamivudine, tenofovir, and/or other renally eliminated drugs. Some examples of drugs that are eliminated by active tubular secretion include, but are not limited to, acyclovir, cidofovir, ganciclovir, valacyclovir, valganciclovir, aminoglycosides (e.g., gentamicin), and high-dose or multiple NSAIDs [see *Warnings and Precautions* (5.3) and *Clinical Pharmacology* (12.3)].

No clinically significant changes in concentration were observed for doravirine when co-administered with the following agents: TDF, lamivudine, elbasvir and grazoprevir, ledipasvir and sofosbuvir, ritonavir, ketoconazole, aluminum hydroxide/magnesium hydroxide/simethicone containing antacid, pantoprazole, or methadone [see *Clinical Pharmacology* (12.3)].

No clinically significant changes in concentration were observed for tenofovir when co-administered with tacrolimus or entecavir [see *Clinical Pharmacology* (12.3)].

7.3 Effect of DELSTRIGO on Other Drugs

No clinically significant changes in concentration were observed for the following agents when co-administered with doravirine: lamivudine, TDF, elbasvir and grazoprevir, ledipasvir and sofosbuvir, atorvastatin, an oral contraceptive containing ethinyl estradiol and levonorgestrel, metformin, methadone, or midazolam.

No clinically significant drug interactions have been observed between TDF and the following medications: entecavir, methadone, oral contraceptives, sofosbuvir, or tacrolimus in studies conducted in healthy participants.

Lamivudine is not significantly metabolized by CYP enzymes nor does it inhibit or induce this enzyme system; therefore, it is unlikely that clinically significant drug interactions will occur through these pathways [see *Clinical Pharmacology* (12.3)].

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Exposure Registry

There is a pregnancy exposure registry that monitors pregnancy outcomes in individuals exposed to DELSTRIGO during pregnancy. Healthcare providers are encouraged to register patients by calling the Antiretroviral Pregnancy Registry (APR) at 1-800-258-4263.

Risk Summary

There is insufficient prospective pregnancy data from the APR to adequately assess the risk of birth defects and miscarriage. Doravirine use in individuals during pregnancy has not been evaluated; however, lamivudine and TDF use during pregnancy has been evaluated in a limited number of individuals reported to the APR. Available data from the APR show no difference in the overall risk of major birth defects for lamivudine and TDF compared with the background rate for major birth defects of 2.7% in the U.S. reference population of the Metropolitan Atlanta Congenital Defects Program (MACDP) (see *Data*). The rate of miscarriage is not reported in the APR. The estimated background rate of miscarriage in the clinically recognized pregnancies in the U.S. general population is 15-20%. Methodological limitations of the APR include the use of MACDP as the external comparator group. The MACDP population is not disease-specific, evaluates individuals and infants from the limited geographic area, and does not include outcomes for births that occurred at less than 20 weeks gestation.

In animal reproduction studies, oral administration of lamivudine to pregnant rabbits during organogenesis resulted in embryoletality at systemic exposure (AUC) similar to the recommended clinical dose; however, no adverse development effects were observed with oral administration of lamivudine to pregnant rats during organogenesis at plasma concentrations (C_{max}) 35 times the recommended clinical dose.

No adverse developmental effects were observed when doravirine and TDF were administered separately at doses/exposures ≥ 8 (doravirine) and ≥ 14 (TDF) times those of the recommended human dose (RHD) of DELSTRIGO (see *Data*).

Data

Human Data

Lamivudine: The APR has received a total of over 13,000 prospective reports with follow-up data of possible exposure to lamivudine-containing regimens; over 5,900 reports in the first trimester; over 5,600 reports in

the second trimester; and over 1,800 reports in the third trimester. Birth defects occurred in 170 of 5,472 (3.1%, 95% CI: 2.7% to 3.6%) live births for lamivudine-containing regimens (first trimester exposure); and 218 of 7,513 (2.9%, 95% CI: 2.5% to 3.3%) live births for lamivudine-containing regimens (second/third trimester exposure). Among pregnant mothers in the U.S. reference population, the background rate of birth defects is 2.7%. There was no association between lamivudine and overall birth defects observed in the APR.

TDF: The APR has received a total of over 7,000 prospective reports with follow-up data of possible exposure to tenofovir disoproxil-containing regimens; over 5,100 reports in the first trimester; over 1,300 reports in the second trimester; and over 600 reports in the third trimester. Birth defects occurred in 113 of 4,576 (2.5%, 95% CI: 2.0% to 3.0%) live births for TDF-containing regimens (first trimester exposure); and 51 of 1,965 (2.6%, 95% CI: 1.9% to 3.4%) live births for TDF-containing regimens (second/third trimester exposure). Among pregnant mothers in the U.S. reference population, the background rate of birth defects is 2.7%. There was no association between tenofovir and overall birth defects observed in the APR.

Animal Data

Doravirine: Doravirine was administered orally to pregnant rabbits (up to 300 mg/kg/day on gestation days (GD) 7 to 20) and rats (up to 450 mg/kg/day on GD 6 to 20 and separately from GD 6 to lactation/postpartum day 20). No significant toxicological effects on embryo-fetal (rats and rabbits) or pre/post-natal (rats) development were observed at exposures (AUC) approximately 9 times (rats) and 8 times (rabbits) the exposure in humans at the RHD. Doravirine was transferred to the fetus through the placenta in embryo-fetal studies, with fetal plasma concentrations of up to 40% (rabbits) and 52% (rats) that of maternal concentrations observed on GD 20.

Lamivudine: Lamivudine was administered orally to pregnant rats (at 90, 600, and 4,000 mg per kg per day) and rabbits (at 90, 300, and 1,000 mg per kg per day and at 15, 40, and 90 mg per kg per day) during organogenesis (on GD 7 through 16 [rat] and 8 through 20 [rabbit]). No evidence of fetal malformations due to lamivudine was observed in rats and rabbits at doses producing plasma concentrations (C_{max}) approximately 35 times higher than human exposure at the recommended daily dose. Evidence of early embryolethality was seen in the rabbit at system exposures (AUC) similar to those observed in humans, but there was no indication of this effect in the rat at plasma concentrations (C_{max}) 35 times higher than human exposure at the recommended daily dose. Studies in pregnant rats showed that lamivudine is transferred to the fetus through the placenta. In the fertility/pre- and postnatal development study in rats, lamivudine was administered orally at doses of 180, 900, and 4,000 mg per kg per day (from prior to mating through postnatal Day 20). In the study, development of the offspring, including fertility and reproductive performance, was not affected by maternal administration of lamivudine.

TDF: Reproduction studies have been performed in rats and rabbits at doses up to 14 and 19 times the human dose based on body surface area comparisons and revealed no evidence of harm to the fetus.

8.2 Lactation

Risk Summary

Based on limited published data, both lamivudine and tenofovir are present in human milk. It is unknown whether doravirine is present in human milk, but doravirine is present in the milk of lactating rats (*see Data*). It is not known whether DELSTRIGO or the components of DELSTRIGO affects human milk production, or has effects on the breastfed infant. Potential risks of breastfeeding include: (1) HIV-1 transmission (in HIV-1-negative infants), (2) developing viral resistance (in HIV-1-positive infants), and (3) serious adverse reactions in a breastfed infant similar to those seen in adults.

Data

Doravirine: Doravirine was excreted into the milk of lactating rats following oral administration (450 mg/kg/day) from GD 6 to lactation day 14, with milk concentrations approximately 1.5 times that of maternal plasma concentrations observed 2 hours post dose on lactation day 14.

8.4 Pediatric Use

The safety and efficacy of DELSTRIGO for the treatment of HIV-1 infection have been established in pediatric patients weighing at least 35 kg [see *Indications and Usage (1)* and *Dosage and Administration (2.2)*].

Use of DELSTRIGO in this group is supported by evidence from adequate and well-controlled trials in adults with additional pharmacokinetic, safety, and efficacy data from an open-label trial in virologically-suppressed or treatment-naïve pediatric participants 12 to less than 18 years of age. The safety and efficacy of DELSTRIGO in these pediatric participants were similar to that in adults, and there was no clinically significant difference in exposure for the components of DELSTRIGO. [see *Adverse Reactions (6.1)*, *Clinical Pharmacology (12.3)*, and *Clinical Studies (14.3)*].

Safety and efficacy of DELSTRIGO in pediatric patients weighing less than 35 kg have not been established.

8.5 Geriatric Use

Clinical trials of doravirine, lamivudine, or TDF did not include sufficient numbers of participants aged 65 years and over to determine whether they respond differently from younger participants. In general, caution should be exercised in the administration of DELSTRIGO in elderly patients reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy [see *Clinical Pharmacology (12.3)*].

8.6 Renal Impairment

Because DELSTRIGO is a fixed-dose combination tablet and the dosage of lamivudine and TDF, both components of DELSTRIGO, cannot be altered, DELSTRIGO is not recommended in patients with estimated creatinine clearance less than 50 mL/min [see *Warnings and Precautions (5.3)* and *Clinical Pharmacology (12.3)*].

8.7 Hepatic Impairment

No dosage adjustment of DELSTRIGO is required in patients with mild (Child-Pugh Class A) or moderate (Child-Pugh Class B) hepatic impairment. DELSTRIGO has not been studied in patients with severe hepatic impairment (Child-Pugh Class C) [see *Clinical Pharmacology (12.3)*].

10 OVERDOSAGE

No data are available on overdose of DELSTRIGO in patients and there is no known specific treatment for overdose with DELSTRIGO. If overdose occurs, the patient should be monitored and standard supportive treatment applied as required.

Doravirine: There is no known specific treatment for overdose with doravirine.

Lamivudine: Because a negligible amount of lamivudine was removed via (4-hour) hemodialysis, continuous ambulatory peritoneal dialysis, and automated peritoneal dialysis, it is not known if continuous hemodialysis would provide clinical benefit in a lamivudine overdose event.

TDF: TDF is efficiently removed by hemodialysis with an extraction coefficient of approximately 54%. Following a single 300 mg dose of TDF, a 4-hour hemodialysis session removed approximately 10% of the administered tenofovir dose.

11 DESCRIPTION

DELSTRIGO is a fixed-dose combination, film-coated tablet, containing doravirine, lamivudine, and TDF for oral administration.

Doravirine is an HIV-1 non-nucleoside reverse transcriptase inhibitor (NNRTI).

Lamivudine is the (-)enantiomer of a dideoxy analogue of cytidine and is an HIV-1 nucleoside analogue reverse transcriptase inhibitor.

TDF (a prodrug of tenofovir) is a fumaric acid salt of the bis-isopropoxycarbonyloxymethyl ester derivative of tenofovir. *In vivo* TDF is converted to tenofovir, an acyclic nucleoside phosphonate (nucleotide) analog of adenosine 5'-monophosphate. Tenofovir is an HIV-1 reverse transcriptase inhibitor.


Each tablet contains 100 mg of doravirine, 300 mg of lamivudine, and 300 mg of TDF (equivalent to 245 mg of tenofovir disoproxil) as active ingredients. The tablets include the following inactive ingredients: colloidal silicon dioxide, croscarmellose sodium, hypromellose acetate succinate, magnesium stearate, microcrystalline cellulose, and sodium stearyl fumarate. The tablets are film coated with a coating material containing the following inactive ingredients: hypromellose, iron oxide yellow, lactose monohydrate, titanium dioxide, and triacetin. The coated tablets are polished with carnauba wax.

Doravirine:

The chemical name for doravirine is 3-chloro-5-[[1-[(4,5-dihydro-4-methyl-5-oxo-1*H*-1,2,4-triazol-3-yl)methyl]-1,2-dihydro-2-oxo-4-(trifluoromethyl)-3-pyridinyl]oxy]benzonitrile.

It has a molecular formula of $C_{17}H_{11}ClF_3N_5O_3$ and a molecular weight of 425.75.

It has the following structural formula:




Doravirine is practically insoluble in water.

Lamivudine:

The chemical name for lamivudine is (-)-1-[(2*R*,5*S*)-2-(hydroxymethyl)-1,3-oxathiolan-5-yl]-cytosine.

It has a molecular formula of $C_8H_{11}N_3O_3S$ and a molecular weight of 229.26.

It has the following structural formula:




Lamivudine is soluble in water.

TDF:

The chemical name for TDF is 9-[(R)-2-[[bis[[[(isopropoxycarbonyl)oxy]methoxy] phosphinyl]-methoxy]propyl]adenine fumarate (1:1).

It has a molecular formula of $C_{19}H_{30}N_5O_{10} \cdot P \cdot C_4H_4O_4$ and a molecular weight of 635.52.

It has the following structural formula:



TDF is slightly soluble in water.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

DELSTRIGO is a fixed-dose combination of the antiretroviral drugs doravirine, lamivudine, and TDF [see *Microbiology* (12.4)].

12.2 Pharmacodynamics

In a Phase 2 trial evaluating doravirine over a dose range of 0.25 to 2 times the recommended dose of doravirine in DELSTRIGO (in combination with FTC/TDF) in participants living with HIV with no antiretroviral treatment history, no exposure-response relationship for efficacy was identified for doravirine.

Cardiac Electrophysiology

At a doravirine dose of 1200 mg, which provides approximately 4 times the peak concentration observed following the recommended dose of doravirine in DELSTRIGO does not prolong the QT interval to any clinically relevant extent.

12.3 Pharmacokinetics

Single-dose administration of one DELSTRIGO tablet to healthy participants provided comparable exposures of doravirine, lamivudine, and tenofovir to administration of doravirine tablets (100 mg) plus lamivudine tablets (300 mg) plus TDF tablets (300 mg). Doravirine pharmacokinetics are similar in healthy participants and participants living with HIV. Pharmacokinetic properties of the components of DELSTRIGO are provided in Table 7.

Table 7: Pharmacokinetic Properties of the Components of DELSTRIGO

Parameter	Doravirine	Lamivudine	Tenofovir
General			
<i>Steady State Exposure*</i>			
AUC ₀₋₂₄ (mcg•h/mL)	16.1 (29) [†]	8.87 ± 1.83 [‡]	2.29 ± 0.69 [§]
C _{max}	0.962 (19) [†]	2.04 ± 0.54 [‡]	0.30 ± 0.09 [§]

(mcg/mL)			
C ₂₄ (mcg/mL)	0.396 (63) [†]	NA	NA
Absorption			
Absolute Bioavailability	64%	86%	25%
T _{max} (h)	2	NA	1
<i>Effect of Food</i> [¶]			
AUC Ratio	1.10 (1.01, 1.20)	0.93 (0.84, 1.03)	1.27 (1.17, 1.37)
C _{max} Ratio	0.95 (0.80, 1.12)	0.81 (0.65, 1.01)	0.88 (0.74, 1.04)
C ₂₄ Ratio	1.26 (1.13, 1.41)	NA	NA
Distribution			
V _{dss} [#]	60.5 L	1.3 L/kg	1.3 L/kg
Plasma Protein Binding	76%	< 36%	<0.7%
Elimination			
t _{1/2} (h)	15	5-7	17
CL/F (mL/ min)*	106 (35.2)	398.5 ± 69.1	1,043.7 ± 115.4
CL _{renal} (mL/ min)*	9.3 (18.6)	199.7 ± 56.9	243.5 ± 33.3
<i>Metabolism</i>			
Primary Pathway(s)	CYP3A	Minor	No CYP Metabolism
<i>Excretion</i>			
Major route of elimination	Metabolism	Glomerular filtration and active tubular secretion	Glomerular filtration and active tubular secretion
Urine (unchanged)	6%	71%	70-80%
Biliary/Fecal (unchanged)	Minor	NA	NA
<p>*Presented as geometric mean (%CV: geometric coefficient of variation) or mean ± SD.</p> <p>[†]Doravirine 100 mg once daily to participants living with HIV.</p> <p>[‡]Lamivudine 300 mg once daily for 7 days to 60 healthy participants.</p> <p>[§]Single 300 mg dose of TDF to participants living with HIV in the fasted state.</p> <p>[¶]Geometric mean ratio [high-fat meal/fasting] and (90% confidence interval) for PK parameters. High fat meal is approximately 1000 kcal, 50% fat. The effect of food is not clinically relevant.</p> <p>[#]Based on IV dose.</p> <p>Abbreviations: NA=not available; AUC=area under the time concentration curve; C_{max}=maximum concentration; C₂₄=concentration at 24 hours; T_{max}=time to C_{max}; V_{dss}=apparent volume of distribution at steady state; t_{1/2}=elimination half-life; CL/F=apparent clearance; CL_{renal} = renal clearance</p>			

Specific Populations

In adults, no clinically significant differences in the pharmacokinetics of certain DELSTRIGO components were observed based on age ≥65 years (for doravirine), sex (for doravirine, lamivudine, tenofovir), and race/ethnicity (for doravirine, lamivudine). The effects of age (≥65 years) on the pharmacokinetics of lamivudine and tenofovir, and the effect of race on the pharmacokinetics of tenofovir are unknown.

Patients with Renal Impairment

Doravirine: No clinically significant difference in the pharmacokinetics of doravirine were observed in participants with mild to severe renal impairment (creatinine clearance (CL_{cr}) >15 mL/min, estimated by Cockcroft-Gault). Doravirine has not been studied in patients with end-stage renal disease or in patients undergoing dialysis.

Lamivudine: The AUC_∞, C_{max}, and half-life of lamivudine increased and CL/F decreased to a clinically significant extent with diminishing renal function (CL_{cr} 111 to < 10 mL/min).

TDF: A clinically significant increase in the C_{max} and AUC of tenofovir was observed in participants with $CL_{cr} < 50$ mL/min or with end stage renal disease requiring dialysis [see *Warnings and Precautions* (5.3) and *Use in Specific Populations* (8.6)].

Patients with Hepatic Impairment

Doravirine: No clinically significant difference in the pharmacokinetics of doravirine was observed in participants with moderate hepatic impairment (Child-Pugh score B) compared to participants without hepatic impairment. Doravirine has not been studied in participants with severe hepatic impairment (Child-Pugh score C).

Lamivudine: No clinically significant differences in lamivudine pharmacokinetics were observed with diminishing hepatic function. Safety and efficacy of lamivudine have not been established in the presence of decompensated liver disease.

TDF: No clinically significant differences in tenofovir pharmacokinetics were observed between participants with any degree of hepatic impairment and unimpaired participants.

Pediatric Patients

Mean doravirine exposures were similar in 54 pediatric participants aged 12 to less than 18 years and weighing at least 35 kg who received doravirine or DELSTRIGO in IMPAACT 2014 (Protocol 027) relative to adults following administration of doravirine or DELSTRIGO. Exposures of lamivudine and tenofovir in pediatric participants following the administration of DELSTRIGO were similar to those in adults following administration of lamivudine and tenofovir (Table 8). For pediatric participants weighing ≥ 35 kg and < 45 kg who receive doravirine 100 mg or DELSTRIGO, the population pharmacokinetic model-predicted mean C_{24} of doravirine was comparable to that achieved in adults, whereas mean AUC_{0-24} and C_{max} of doravirine were 25% and 36% higher than adult values, respectively. However, the predicted AUC_{0-24} and C_{max} increases are not considered clinically significant.

Table 8: Steady State Pharmacokinetics for Doravirine, Lamivudine, and Tenofovir Following Administration of Doravirine or DELSTRIGO in Pediatric Participants Living with HIV Aged 12 to Less than 18 Years and Weighing at Least 35 kg

Parameter*	Doravirine [†]	Lamivudine [‡]	Tenofovir [‡]
AUC_{0-24} (mcg•h/mL)	16.4 (24)	11.3 (28)	2.55 (14)
C_{max} (mcg/mL)	1.03 (16)	2.1 (24)	0.293 (37)
C_{24} (mcg/mL)	0.379 (42)	NA	NA
*Presented as geometric mean (%CV: geometric coefficient of variation) [†] From population PK analysis (n=53 weighing ≥ 45 kg, n=1 weighing ≥ 35 kg to < 45 kg) [‡] From intensive PK analysis (n=10) Abbreviations: NA=not applicable; AUC=area under the time concentration curve; C_{max} =maximum concentration; C_{24} =concentration at 24 hours			

Drug Interaction Studies

DELSTRIGO is a complete regimen for the treatment of HIV-1 infection; therefore, DELSTRIGO is not recommended to be administered with other HIV-1 antiretroviral medications. Information regarding potential drug-drug interactions with other antiretroviral medications is not provided.

The drug interaction trials described were conducted with doravirine, lamivudine and/or TDF, as single entities; no drug interaction trials have been conducted using the combination of doravirine, lamivudine, and TDF. No clinically relevant drug interactions were observed between doravirine, lamivudine, and TDF.

Doravirine: Doravirine is primarily metabolized by CYP3A, and drugs that induce or inhibit CYP3A may affect the clearance of doravirine. Co-administration of doravirine and drugs that induce CYP3A may result in decreased plasma concentrations of doravirine. Co-administration of doravirine and drugs that inhibit CYP3A may result in increased plasma concentrations of doravirine.

Doravirine is not likely to have a clinically relevant effect on the exposure of medicinal products metabolized by CYP enzymes. Doravirine did not inhibit major drug metabolizing enzymes *in vitro*, including CYPs 1A2, 2B6, 2C8, 2C9, 2C19, 2D6, 3A4, and UGT1A1 and is not likely to be an inducer of CYP1A2, 2B6, or 3A4. Based on *in vitro* assays, doravirine is not likely to be an inhibitor of OATP1B1, OATP1B3, P-glycoprotein, BSEP, OAT1, OAT3, OCT2, MATE1, and MATE2K. Drug interaction studies were performed with doravirine and other drugs likely to be co-administered or commonly used as probes for pharmacokinetic interactions. The effects of co-administration with other drugs on the exposure (C_{max} , AUC, and C_{24}) of doravirine are summarized in Table 9. A single doravirine 100 mg dose was administered in these studies unless otherwise noted.

Table 9: Drug Interactions: Changes in Pharmacokinetic Parameter Values of Doravirine in the Presence of Co-administered Drug

Co-administered Drug	Regimen of Co-administered Drug	N	Geometric Mean Ratio (90% CI) of Doravirine Pharmacokinetics with/without Co-administered Drug (No Effect=1.00)		
			AUC*	C _{max}	C ₂₄
Azole Antifungal Agents					
ketoconazole [†]	400 mg QD	10	3.06 (2.85, 3.29)	1.25 (1.05, 1.49)	2.75 (2.54, 2.98)
Antimycobacterials					
rifampin	600 mg QD	10	0.12 (0.10, 0.15)	0.43 (0.35, 0.52)	0.03 (0.02, 0.04)
rifabutin	300 mg QD	12	0.50 (0.45, 0.55)	0.99 (0.85, 1.15)	0.32 (0.28, 0.35)
	300 mg QD [‡]	15	1.03 (0.94, 1.14)	0.97 (0.87, 1.08)	0.98 (0.88, 1.10)
HIV Antiviral Agents					
ritonavir ^{†,§}	100 mg BID	8	3.54 (3.04, 4.11)	1.31 (1.17, 1.46)	2.91 (2.33, 3.62)
efavirenz	600 mg QD [¶]	17	0.38 (0.33, 0.45)	0.65 (0.58, 0.73)	0.15 (0.10, 0.23)
	600 mg QD [#]	17	0.68 (0.58, 0.80)	0.86 (0.77, 0.97)	0.50 (0.39, 0.64)

CI = confidence interval; QD = once daily; BID = twice daily

*AUC_{0-∞} for single-dose, AUC₀₋₂₄ for once daily.

[†]Changes in doravirine pharmacokinetic values are not clinically relevant.

[‡]Doravirine 100 mg BID resulted in similar pharmacokinetic values when compared to 100 mg QD without rifabutin.

[§]A single doravirine 50 mg dose (0.5 times the recommended approved dose) was administered.

[¶]The first day following the cessation of efavirenz therapy and initiation of doravirine 100 mg QD.

[#]14 days following the cessation of efavirenz therapy and initiation of doravirine 100 mg QD.

Based on drug interaction studies conducted with doravirine, no clinically significant drug interactions have been observed following the co-administration of doravirine and the following drugs: dolutegravir, TDF, lamivudine, elbasvir and grazoprevir, ledipasvir and sofosbuvir, ketoconazole, ritonavir, aluminum hydroxide/magnesium hydroxide/simethicone containing antacid, pantoprazole, atorvastatin, an oral contraceptive containing ethinyl estradiol and levonorgestrel, metformin, methadone, and midazolam.

Lamivudine:

Trimethoprim/Sulfamethoxazole: Co-administration of TMP/SMX with lamivudine resulted in an increase of 43% ±23% (mean ±SD) in lamivudine AUC_∞, a decrease of 29% ±13% in lamivudine oral clearance, and a decrease of 30% ±36% in lamivudine renal clearance. The pharmacokinetic properties of TMP and SMX were not altered by co-administration with lamivudine.

Sorbitol (Excipient): Co-administration of lamivudine with a single dose of 3.2 grams, 10.2 grams, or 13.4 grams of sorbitol resulted in dose-dependent decreases of 14%, 32%, and 36% in the AUC_∞; and 28%, 52%, and 55% in the C_{max} of lamivudine, respectively.

TDF:

No clinically significant changes in exposure were observed for tenofovir when co-administered with tacrolimus or entecavir.

No clinically significant changes in exposure were observed for the following drugs when co-administered with tenofovir: tacrolimus, entecavir, methadone, or ethinyl estradiol/norgestimate.

12.4 Microbiology

Mechanism of Action

Doravirine: Doravirine is a pyridinone non-nucleoside reverse transcriptase inhibitor of HIV-1 and inhibits HIV-1 replication by non-competitive inhibition of HIV-1 reverse transcriptase (RT). The inhibitory concentration at 50% (IC₅₀) of doravirine for RNA-dependent DNA polymerization of recombinant wild-type HIV-1 RT in a biochemical assay was 12.2±2.0 nM (n=3). Doravirine does not inhibit the human cellular DNA polymerases α, β, and mitochondrial DNA polymerase γ.

Lamivudine: Lamivudine is a synthetic nucleoside analogue. Intracellularly, lamivudine is phosphorylated to its active 5'-triphosphate metabolite, lamivudine triphosphate (3TC-TP). The principal mode of action of 3TC-TP is inhibition of RT via DNA chain termination after incorporation of the nucleotide analogue. Lamivudine triphosphate (3TC-TP) is a weak inhibitor of mammalian DNA polymerases α, β, and mitochondrial DNA polymerase γ.

TDF: TDF is an acyclic nucleoside phosphonate diester analog of adenosine monophosphate. TDF requires initial diester hydrolysis for conversion to tenofovir and subsequent phosphorylations by cellular enzymes to form tenofovir diphosphate. Tenofovir diphosphate inhibits the activity of HIV-1 RT by competing with the natural substrate deoxyadenosine 5'-triphosphate and, after incorporation into DNA, by DNA chain termination. Tenofovir diphosphate is a weak inhibitor of mammalian DNA polymerases α, β, and mitochondrial DNA polymerase γ.

Antiviral Activity in Cell Culture

Doravirine: Doravirine exhibited an EC₅₀ value of 12.0±4.4 nM against wild-type laboratory strains of HIV-1 when tested in the presence of 100% normal human serum (NHS) using MT4-GFP reporter cells and a median EC₅₀ value for HIV-1 subtype B primary isolates (n=118) of 4.1 nM (range: 1.0 nM-16.0 nM). Doravirine demonstrated antiviral activity against a broad panel of primary HIV-1 isolates (A, A1, AE, AG, B, BF, C, D, G, H) with EC₅₀ values ranging from 1.2 nM to 10.0 nM. The antiviral activity of doravirine was not antagonistic when combined with lamivudine and TDF.

Lamivudine: The antiviral activity of lamivudine against HIV-1 was assessed in a number of cell lines including monocytes and peripheral blood mononuclear cells (PBMCs) using standard susceptibility assays. EC₅₀ values were in the range of 3 to 15,000 nM (1,000 nM = 230 ng per mL). The median EC₅₀ values of lamivudine were 60 nM (range: 20 to 70 nM), 35 nM (range: 30 to 40 nM), 30 nM (range: 20 to 90 nM), 20 nM (range: 3 to 40 nM), 30 nM (range: 1 to 60 nM), 30 nM (range: 20 to 70 nM), 30 nM (range: 3 to 70 nM), and 30 nM (range: 20 to 90 nM) against HIV-1 clades A-G and group O viruses (n = 3 except n = 2 for clade B) respectively. Ribavirin (50,000 nM) used in the treatment of chronic HCV infection decreased the anti-HIV-1 activity of lamivudine by 3.5-fold in MT-4 cells.

TDF: The antiviral activity of tenofovir against laboratory and clinical isolates of HIV-1 was assessed in T lymphoblastoid cell lines, primary monocyte/macrophage cells and peripheral blood lymphocytes. The EC₅₀

values for tenofovir were in the range of 40-8,500 nM. Tenofovir displayed antiviral activity in cell culture against HIV-1 clades A, B, C, D, E, F, G, and O (EC₅₀ values ranged from 500-2,200 nM).

Resistance

In Cell Culture

Doravirine: Doravirine-resistant strains were selected in cell culture starting from wild-type HIV-1 of different origins and subtypes, as well as NNRTI-resistant HIV-1. Observed emergent amino acid substitutions in RT included: V106A, V106I, V106M, V108I, H221Y, F227C, F227I, F227L, F227V, M230I, L234I, P236L, and Y318F. The V106A, V106M, V108I, H221Y, F227C, M230I, P236L, and Y318F substitutions conferred 3.4-fold to 70-fold reductions in susceptibility to doravirine. Y318F in combination with V106A, V106M, V108I, or F227C conferred greater decreases in susceptibility to doravirine than Y318F alone, which conferred a 10-fold reduction in susceptibility to doravirine.

Lamivudine: Lamivudine-resistant variants of HIV-1 have been selected in cell culture and in participants treated with lamivudine. Genotypic analysis showed that substitutions M184I or V cause resistance to lamivudine.

TDF: HIV-1 isolates selected by tenofovir in cell culture expressed a K65R substitution in HIV-1 RT and showed a 2–4 -fold reduction in susceptibility to tenofovir. In addition, a K70E substitution in HIV-1 RT has been selected by tenofovir and results in low-level reduced susceptibility to abacavir, emtricitabine, lamivudine, and tenofovir.

In Clinical Trials

Clinical Trial Results in Adults with No Antiretroviral Treatment History

Doravirine: In the doravirine treatment arm of the DRIVE-AHEAD trial (n=364) through 96 weeks, 10 participants showed the emergence of doravirine resistance-associated substitutions among 24 (42%) participants in the resistance analysis subset (participants with HIV-1 RNA greater than 400 copies per mL at virologic failure or early study discontinuation and having post-baseline resistance samples). Emergent doravirine resistance-associated substitutions in RT included one or more of the following: V90V/G, A98G, V106A, V106I, V106M/T, V108I, E138G, Y188L, H221Y, P225H, P225L, P225P/S, F227C, F227C/R, Y318Y/F, and Y318Y/S. Six of the 10 participants with emergent doravirine resistance-associated substitutions showed doravirine phenotypic resistance and all of them had a greater than 100-fold reduction in doravirine susceptibility (range >103 to >211). The other 4 virologic failures who had only amino acid mixtures of NNRTI resistance substitutions showed doravirine phenotypic fold-changes of less than 2-fold.

In the EFV/FTC/TDF treatment arm of the DRIVE-AHEAD trial (n=364) through Week 96, 15 participants showed the emergence of efavirenz resistance-associated substitutions among 25 (60%) participants in the resistance analysis subset.

Lamivudine and TDF: In a pooled analysis of antiretroviral-naïve participants who received doravirine, lamivudine, and TDF, genotyping was performed on plasma HIV-1 isolates from all participants with HIV-1 RNA greater than 400 copies per mL at confirmed virologic failure or at time of early study drug discontinuation. Genotypic resistance developed in 8 evaluable participants who received DOR/3TC/TDF through Week 96. The resistance-associated substitutions that emerged were RT M41L (n=1), A62A/V (n=1), K65R (n=2), T69T/A (n=1), V75V/I (n=1), and M184V (n=5). In comparison, genotypic resistance to emtricitabine or tenofovir developed in 5 evaluable participants who received EFV/FTC/TDF in DRIVE-AHEAD; emergent resistance-associated substitutions were RT K65R (n=1), D67G/K70E (n=1), L74V/V75M/V118I (n=1), M184I or V (n=5), and K219K/E (n=1).

Clinical Trial Results in Virologically-Suppressed Adults

In the DRIVE-SHIFT clinical trial [see *Clinical Studies* (14.2)], there were 6 participants in the immediate switch group (n=447) and 2 participants in the delayed switch group (n=209) who met the protocol-defined virologic failure criteria (confirmed HIV-1 RNA \geq 50 copies/mL). Two of the 6 virologic failure participants in

the immediate switch group had available resistance data and neither developed detectable genotypic or phenotypic resistance to doravirine, lamivudine, or tenofovir during treatment with DELSTRIGO. One of the two virologic failure participants in the delayed switch group who had available resistance data developed the RT M184M/I substitution and phenotypic resistance to emtricitabine and lamivudine during treatment with their baseline regimen.

Cross-Resistance

No significant cross-resistance has been demonstrated between doravirine-resistant HIV-1 variants and lamivudine/emtricitabine or tenofovir or between lamivudine or tenofovir-resistant variants and doravirine.

Doravirine: Cross-resistance has been observed among NNRTIs. Treatment-emergent doravirine resistance-associated substitutions can confer cross resistance to efavirenz, etravirine, nevirapine, and rilpivirine. Of the 6 virologic failure participants who developed doravirine phenotypic resistance, all had phenotypic resistance to efavirenz and nevirapine, 4 had phenotypic resistance to rilpivirine, and 4 had resistance to etravirine in the Monogram PhenoSense assay. Of the 11 virologic failure participants phenotypically resistant to efavirenz, 2 (18%) had decreased susceptibility to doravirine (18- and 36-fold).

The treatment-emergent doravirine resistance-associated substitution Y318F did not confer reduced susceptibility to efavirenz, etravirine, or rilpivirine.

A panel of 96 diverse clinical isolates containing NNRTI resistance-associated substitutions was evaluated for susceptibility to doravirine. Clinical isolates containing the Y188L substitution alone or in combination with K103N or V106I, V106A in combination with G190A and F227L, or E138K in combination with Y181C and M230L showed greater than 100-fold reduced susceptibility to doravirine.

Lamivudine: Cross-resistance has been observed among NRTIs. The M184I/V lamivudine resistance-associated substitution confers resistance to abacavir, didanosine and emtricitabine. Lamivudine also has reduced susceptibility against the K65R substitution.

TDF: Cross-resistance has been observed among NRTIs. The K65R substitution in HIV-1 RT selected by tenofovir is also selected in some patients living with HIV treated with abacavir or didanosine. HIV-1 isolates with the K65R substitution also showed reduced susceptibility to emtricitabine and lamivudine. Therefore, cross-resistance among these NRTIs may occur in patients whose virus harbors the K65R substitution. The K70E substitution selected clinically by TDF results in reduced susceptibility to abacavir, didanosine, emtricitabine, lamivudine, and tenofovir. HIV-1 isolates from patients (n=20) whose HIV-1 expressed a mean of 3 zidovudine resistance-associated substitutions (M41L, D67N, K70R, L210W, T215Y/F, or K219Q/E/N) showed a 3.1-fold decrease in the susceptibility to tenofovir. Participants whose virus expressed an RT L74V substitution without zidovudine resistance-associated substitutions (n=8) had reduced response to TDF. Limited data are available for patients whose virus expressed a Y115F substitution (n=3), Q151M substitution (n=2), or T69 insertion (n=4) in HIV-1 RT, all of whom had a reduced response in clinical trials.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Doravirine: Doravirine was not carcinogenic in long-term oral carcinogenicity studies in mice and rats at exposures up to 6 and 7 times, respectively, the human exposures at the RHD. A statistically significant incidence of thyroid parafollicular cell adenoma and carcinoma seen only in female rats at the high dose was within the range observed in historical controls.

Lamivudine: Long-term carcinogenicity studies with lamivudine in mice and rats showed no evidence of carcinogenic potential at exposures up to 10 times (mice) and 58 times (rats) the human exposures at the RHD.

TDF: Long-term oral carcinogenicity studies of TDF in mice and rats were carried out at exposures up to approximately 16 times (mice) and 5 times (rats) those observed in humans at the RHD. At the high dose in female mice, liver adenomas were increased at exposures 16 times of that in humans. In rats, the study was negative for carcinogenic findings at exposures up to 5 times that observed in humans at the RHD.

Mutagenesis

Doravirine: Doravirine was not genotoxic in a battery of *in vitro* or *in vivo* assays, including microbial mutagenesis, chromosomal aberration in Chinese hamster ovary cells, and in *in vivo* rat micronucleus assays.

Lamivudine: Lamivudine was mutagenic in an L5178Y mouse lymphoma assay and clastogenic in a cytogenetic assay using cultured human lymphocytes. Lamivudine was not mutagenic in a microbial mutagenicity assay, in an *in vitro* cell transformation assay, in a rat micronucleus test, in a rat bone marrow cytogenetic assay, and in an assay for unscheduled DNA synthesis in rat liver. Lamivudine showed no evidence of *in vivo* genotoxic activity in the rat at oral doses of up to 2,000 mg per kg, producing plasma levels of 35 to 45 times those in humans at the recommended dose for HIV-1 infection.

TDF: TDF was mutagenic in the *in vitro* mouse lymphoma assay and negative in an *in vitro* bacterial mutagenicity test (Ames test). In an *in vivo* mouse micronucleus assay, TDF was negative when administered to male mice.

Impairment of Fertility

Doravirine: There were no effects on fertility, mating performance or early embryonic development when doravirine was administered to rats up to the highest dose tested. Systemic exposures (AUC) to doravirine were approximately 7 times the exposure in humans at the RHD.

Lamivudine: In a study of reproductive performance, lamivudine administered to rats at doses up to 4,000 mg per kg per day, producing plasma levels 47 to 70 times those in humans, revealed no evidence of impaired fertility and no effect on the survival, growth, and development to weaning of the offspring.

TDF: There were no effects on fertility, mating performance or early embryonic development when TDF was administered to male rats at a dose equivalent to 10 times the RHD based on body surface area comparisons for 28 days prior to mating and to female rats for 15 days prior to mating through day 7 of gestation. There was, however, an alteration of the estrous cycle in female rats.

14 CLINICAL STUDIES

14.1 Clinical Trial Results in Adults with No Antiretroviral Treatment History

The efficacy of DELSTRIGO is based on the analyses of 96-week data from a randomized, multicenter, double-blind, active controlled Phase 3 trial (DRIVE-AHEAD, NCT02403674) in participants living with HIV with no antiretroviral treatment history (n=728).

Participants were randomized and received at least 1 dose of either DELSTRIGO or EFV 600 mg/FTC 200 mg/TDF 300 mg once daily. At baseline, the median age of participants was 31 years, 15% were female, 52% were Non-White, 3% had hepatitis B or C coinfection, 14% had a history of AIDS, 21% had HIV-1 RNA greater than 100,000 copies/mL, and 88% had CD4+ T-cell count greater than 200 cells/mm³; these characteristics were similar between treatment groups. Week 96 outcomes for DRIVE-AHEAD are provided in Table 10.

Mean CD4+ T-cell counts in the DELSTRIGO and EFV/FTC/TDF groups increased from baseline by 238 and 223 cells/mm³, respectively.

Table 10: Virologic Outcome in DRIVE-AHEAD at Week 96 in HIV-1 Adult Participants with No Antiretroviral Treatment History

Outcome	DELSTRIGO Once Daily N=364	EFV/FTC/TDF Once Daily N=364
HIV-1 RNA <50 copies/mL	77%	74%
Treatment Difference (95% CI) *	3.8% (-2.4%, 10.0%)	
HIV-1 RNA ≥ 50 copies/mL[†]	15%	12%
No Virologic Data at Week 96 Window	7%	14%
Discontinued study due to AE or Death [‡]	3%	8%
Discontinued study for Other Reasons [§]	4%	5%
On study but missing data in window	1%	1%
Proportion (%) of Participants With HIV-1 RNA <50 copies/mL at Week 96 by Baseline and Demographic Category		
Gender		
Male	78% (N = 305)	73% (N = 311)
Female	75% (N = 59)	75% (N = 53)
Race		
White	80% (N = 176)	74% (N = 170)
Non-White	76% (N = 188)	74% (N = 194)
Ethnicity[¶]		
Hispanic or Latino	81% (N = 126)	77% (N = 119)
Not Hispanic or Latino	76% (N = 238)	72% (N = 239)
Baseline HIV-1 RNA (copies/mL)		
≤100,000 copies/mL	80% (N = 291)	77% (N = 282)
>100,000 copies/mL	67% (N = 73)	62% (N = 82)
CD4+ T-cell Count (cells/mm³)		
≤200 cells/mm ³	59% (N = 44)	70% (N = 46)
>200 cells/mm ³	80% (N = 320)	74% (N = 318)
Viral Subtype[¶]		
Subtype B	80% (N = 232)	72% (N = 253)
Subtype Non-B	73% (N = 130)	77% (N = 111)
<p>*The 95% CI for the treatment difference was calculated using stratum-adjusted Mantel-Haenszel method.</p> <p>[†]Includes participants who discontinued study drug or study before Week 96 for lack or loss of efficacy and participants with HIV-1 RNA equal to or above 50 copies/mL in the Week 96 window.</p> <p>[‡]Includes participants who discontinued because of adverse event (AE) or death if this resulted in no virologic data in the Week 96 window.</p> <p>[§]Other reasons include: lost to follow-up, non-compliance with study drug, physician decision, pregnancy, protocol deviation, screen failure, withdrawal by participant.</p> <p>[¶]Does not include participants whose ethnicity or viral subtypes were unknown.</p>		

14.2 Clinical Trial Results in Virologically-Suppressed Adults

The efficacy of switching from a baseline regimen consisting of two NRTIs in combination with a PI plus either ritonavir or cobicistat, or elvitegravir plus cobicistat, or an NNRTI to DELSTRIGO was evaluated in a randomized, open-label trial (DRIVE-SHIFT, NCT02397096), in virologically-suppressed adults living with HIV. Participants must have been virologically-suppressed (HIV-1 RNA <50 copies/mL) on their baseline

regimen for at least 6 months prior to trial entry, with no history of virologic failure. Participants were randomized to either switch to DELSTRIGO at baseline [n = 447, Immediate Switch Group (ISG)], or stay on their baseline regimen until Week 24, at which point they switched to DELSTRIGO [n = 223, Delayed Switch Group (DSG)].

At baseline, the median age of participants was 43 years, 16% were female, and 24% were Non-White, 21% were of Hispanic or Latino ethnicity, 3% had hepatitis B and/or C virus co-infection, 17% had a history of AIDS, 96% had CD4+ T-cell count greater than or equal to 200 cells/mm³, 70% were on a regimen containing a PI plus ritonavir, 24% were on a regimen containing an NNRTI, 6% were on a regimen containing elvitegravir plus cobicistat, and 1% were on a regimen containing a PI plus cobicistat; these characteristics were similar between treatment groups.

Virologic outcome results are shown in Table 11.

Table 11: Virologic Outcomes in DRIVE-SHIFT in HIV-1 Virologically-Suppressed Participants Who Switched to DELSTRIGO

Outcome	DELSTRIGO Once Daily ISG Week 48 N=447	Baseline Regimen DSG Week 24 N=223
HIV-1 RNA \geq 50 copies/mL*	2%	1%
ISG-DSG, Difference (95% CI) ^{†,‡}	0.7% (-1.3%, 2.6%)	
HIV-1 RNA <50 copies/mL	91%	95%
No Virologic Data Within the Time Window	8%	4%
Discontinued study due to AE or Death [§]	3%	<1%
Discontinued study for Other Reasons [¶]	4%	4%
On study but missing data in window	0	0
Proportion (%) of Participants With HIV-1 RNA <50 copies/mL by Baseline and Demographic Category		
Age (years)		
<50	90% (N = 320)	95% (N = 157)
\geq 50	94% (N = 127)	94% (N = 66)
Gender		
Male	91% (N = 372)	94% (N = 194)
Female	91% (N = 75)	100% (N = 29)
Race		
White	90% (N = 344)	95% (N = 168)
Non-White	93% (N = 103)	93% (N = 55)
Ethnicity		
Hispanic or Latino	88% (N = 99)	91% (N = 45)
Not Hispanic or Latino	91% (N = 341)	95% (N = 175)
CD4+ T-cell Count (cells/mm³)		
<200 cells/mm ³	85% (N = 13)	75% (N = 4)
\geq 200 cells/mm ³	91% (N = 426)	95% (N = 216)
Baseline Regimen[#]		
PI plus either ritonavir or cobicistat	90% (N = 316)	94% (N = 156)

elvitegravir plus cobicistat or NNRTI	93% (N = 131)	96% (N = 67)
<p>*Includes participants who discontinued study drug or study before Week 48 for ISG or before Week 24 for DSG for lack or loss of efficacy and participants with HIV-1 RNA ≥ 50 copies/mL in the Week 48 window for ISG and in the Week 24 window for DSG</p> <p>†The 95% CI for the treatment difference was calculated using stratum-adjusted Mantel-Haenszel method.</p> <p>‡Assessed using a non-inferiority margin of 4%.</p> <p>§Includes participants who discontinued because of adverse event (AE) or death if this resulted in no virologic data on treatment during the specified window.</p> <p>¶Other reasons include: lost to follow-up, non-compliance with study drug, physician decision, protocol deviation, withdrawal by participant.</p> <p>#Baseline Regimen = PI plus either ritonavir or cobicistat (specifically atazanavir, darunavir, or lopinavir), or elvitegravir plus cobicistat, or NNRTI (specifically efavirenz, nevirapine, or rilpivirine), each administered with two NRTIs.</p>		

14.3 Clinical Trial Results in Pediatric Participants

The efficacy of DELSTRIGO was evaluated in cohort 2 of an open-label, single-arm 2-cohort trial in pediatric participants 12 to less than 18 years of age living with HIV (IMPAACT 2014 (Protocol 027), NCT03332095). In cohort 1, virologically-suppressed participants (n=9) received a single 100 mg dose of doravirine followed by intensive PK sampling. In cohort 2, virologically-suppressed participants (n=43) were switched to DELSTRIGO and treatment-naïve participants (n=2) were started on DELSTRIGO.

In cohort 2, at baseline the median age of participants was 15 years (range: 12 to 17), the median weight was 52 kg (range: 45 to 80), 58% were female, 78% were Asian and 22% were Black, and the median CD4+ T-cell count was 713 cells per mm³ (range 84 to 1397). After switching to DELSTRIGO, 95% (41/43) of virologically-suppressed participants remained suppressed (HIV-1 RNA <50 copies/mL) at Week 24. One of the two treatment-naïve participants achieved HIV-1 RNA <50 copies/mL at Week 24. The other treatment-naïve participant met the protocol-defined virologic failure criteria (defined as 2 consecutive plasma HIV-1 RNA test results ≥ 200 copies/mL at or after Week 24) and was evaluated for the development of resistance; no emergence of genotypic or phenotypic resistance to doravirine, lamivudine, or tenofovir was detected.

16 HOW SUPPLIED/STORAGE AND HANDLING

Each DELSTRIGO tablet contains 100 mg of doravirine, 300 mg of lamivudine, and 300 mg of tenofovir disoproxil fumarate (equivalent to 245 mg of tenofovir disoproxil), is yellow, oval-shaped, film-coated, and is debossed with the corporate logo and 776 on one side and plain on the other side. Each bottle contains 30 tablets (NDC 0006-5007-01) and silica gel desiccants, and is closed with a child-resistant closure.

Store DELSTRIGO in the original bottle. Keep the bottle tightly closed to protect from moisture. Do not remove the desiccants.

Store DELSTRIGO at 20°C to 25°C (68°F to 77°F); excursions permitted to 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature].

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Patient Information).

Severe Skin Reactions

Inform patients that severe skin reactions including Stevens-Johnson syndrome (SJS)/toxic epidermal necrolysis (TEN) have been reported with DELSTRIGO. Advise patients to immediately contact their healthcare provider if they develop a rash. Instruct patients to immediately stop taking DELSTRIGO and seek medical attention if a painful rash with mucosal involvement develops [see *Warnings and Precautions* (5.1)].

Severe Acute Exacerbation of Hepatitis B in people with concomitant HIV-1 and HBV

Inform patients that severe acute exacerbations of hepatitis B have been reported in people with concomitant HIV-1 and HBV who have discontinued lamivudine or TDF and may occur with discontinuation of DELSTRIGO [see *Warnings and Precautions* (5.2)]. Advise patients not to discontinue DELSTRIGO without first informing their healthcare provider.

Drug Interactions

Inform patients that DELSTRIGO may interact with certain other drugs; therefore, advise patients to report to their healthcare provider the use of any other prescription or nonprescription medication or herbal products, including St. John's wort [see *Contraindications* (4), *Warnings and Precautions* (5.4), and *Drug Interactions* (7)].

For patients concomitantly receiving rifabutin, take one tablet of doravirine (PIFELTRO) 100 mg approximately 12 hours after the dose of DELSTRIGO [see *Dosage and Administration* (2.4)].

New Onset or Worsening Renal Impairment

Inform patients that renal impairment, including cases of acute renal failure and Fanconi syndrome, has been reported in association with the use of TDF. Advise patients to avoid DELSTRIGO with concurrent or recent use of a nephrotoxic agent (e.g., high-dose or multiple NSAIDs) [see *Warnings and Precautions* (5.3)].

Bone Loss and Mineralization Defects

Inform patients that decreases in bone mineral density have been observed with the use of TDF, a component of DELSTRIGO. Assessment of bone mineral density (BMD) should be considered in patients who have a history of pathologic bone fracture or other risk factors for osteoporosis or bone loss [see *Warnings and Precautions* (5.5)].

Immune Reconstitution Syndrome

Inform patients that in some patients with advanced HIV infection (AIDS), signs and symptoms of inflammation from previous infections may occur soon after anti-HIV treatment is started. It is believed that these symptoms are due to an improvement in the body's immune response, enabling the body to fight infections that may have been present with no obvious symptoms. Advise patients to inform their healthcare provider immediately of any symptoms of infection [see *Warnings and Precautions* (5.6)].

Dosing Instructions

Advise patients to take DELSTRIGO every day at a regularly scheduled time with or without food. Inform patients that it is important not to miss or skip doses as it can result in development of resistance. If a patient forgets to take DELSTRIGO, tell the patient to take the missed dose right away, unless it is almost time for the next dose. Advise the patient not to take 2 doses at one time and to take the next dose at the regularly scheduled time.

Pregnancy Registry

Inform patients that there is an antiretroviral pregnancy registry to monitor fetal outcomes of pregnant individuals exposed to DELSTRIGO [see *Use in Specific Populations* (8.1)].

Lactation

Inform individuals with HIV-1 infection that the potential risks of breastfeeding include: (1) HIV-1 transmission (in HIV-1-negative infants), (2) developing viral resistance (in HIV-1-positive infants), and (3) serious adverse reactions in a breastfed infant similar to those seen in adults [see *Use in Specific Populations* (8.2)].

Manufactured for: Merck Sharp & Dohme LLC
Rahway, NJ 07065, USA

For patent information: www.msd.com/research/patent

The trademarks depicted herein are owned by their respective companies.

Copyright © 2018-2025 Merck & Co., Inc., Rahway, NJ, USA, and its affiliates.
All rights reserved.

uspi-mk1439a-t-2506r009