

**INSTRUCTIONS FOR USE**  
**SINGULAIR® (SING-u-lair)**  
**(montelukast sodium)**  
**oral granules**

This Instructions for Use contains information on how to use SINGULAIR oral granules.

**Important Information:**


- Before giving a dose of SINGULAIR oral granules, read this Instructions for Use to be sure you prepare and give the oral granules correctly.
- Give SINGULAIR oral granules to your child exactly as instructed by your healthcare provider.
- Stop giving SINGULAIR and tell your healthcare provider right away if your child has any unusual changes in behavior or thinking.
- Continue to give your child their asthma medicines as prescribed, unless your healthcare provider tells you to change how you give these medicines.
- You can give SINGULAIR oral granules with food or without food.

**How can I give SINGULAIR oral granules to my child?**

- **Do not** open the packet until ready to use.
- There are different ways you can give SINGULAIR 4-mg oral granules. You should choose the best method for your child:
  - right into the mouth
  - dissolved in 1 teaspoonful (5 mL) of cold or room temperature baby formula or breast milk
  - mixed with 1 spoonful of one of the following soft foods at cold or room temperature: applesauce, mashed carrots, rice, or ice cream.
- Give the child all of the mixture within 15 minutes.
- **Do not store any leftover SINGULAIR mixture (oral granules mixed with food, baby formula, or breast milk) for use at a later time.** Throw away any unused portion.
- **Do not mix SINGULAIR oral granules with any liquid drink other than baby formula or breast milk.** Your child may drink other liquids after swallowing the mixture.

**How should I store SINGULAIR?**

- Store SINGULAIR at room temperature between 68°F to 77°F (20°C to 25°C).
- Keep SINGULAIR in the package it comes in.
- Keep SINGULAIR in a dry place and keep it away from light.
- **Keep SINGULAIR and all medicines out of the reach of children.**

 Dist. by: Merck Sharp & Dohme Corp., a subsidiary of  
**MERCK & CO., INC.**, Whitehouse Station, NJ 08889, USA

For patent information: [www.merck.com/product/patent/home.html](http://www.merck.com/product/patent/home.html)  
Copyright © 1998-2020 Merck Sharp & Dohme Corp., a subsidiary of **Merck & Co., Inc.**  
All rights reserved.  
usifu-mk0476-og-2004r001  
For more information, call the Merck National Service Center at 1-800-NSC-Merck (1-800-672-6372)

This Instructions for Use has been approved by the U.S. Food and Drug Administration

Approved: April 2020